

# CHIPPEWA COUNTY FOREST COMPREHENSIVE LAND USE PLAN

## TABLE OF CONTENTS

### CHAPTER 600

#### PROTECTION

<u>Section</u>	<u>Subject</u>	<u>Page</u>
<b>600</b>	<b>PROTECTION.....</b>	<b>3</b>
<b>605</b>	<b>FIRE CONTROL.....</b>	<b>3</b>
605.1	COOPERATION WITH THE DEPARTMENT OF NATURAL RESOURCES.....	3
605.1.1	Personnel.....	4
605.1.2	Equipment.....	4
605.1.3	Fire Detection.....	4
605.1.4	Forest Fire Prevention.....	4
605.2	DEBRIS BURNING.....	4
605.3	CAMPING FIRES.....	5
605.4	PRESCRIBED BURINING.....	5
605.5	COUNTY FOREST FIRE HAZARD AREAS.....	5
<b>610</b>	<b>CONTROL OF FOREST PESTS &amp; PATHOGENS.....</b>	<b>5</b>
610.1	DETECTION.....	5
610.2	PEST SURVEYS.....	5
610.3	SPECIFIC PESTS AND PATHOGENS OF CONCERN.....	6
610.3.1	Specific Pests of Interest.....	6
610.3.1.1	Gypsy Moth .....	6
610.3.1.2	Oak Wilt.....	6
610.3.1.3	Forest Tent Caterpillar.....	7
610.3.1.4	Two-Lined Chestnut Borer.....	7
610.3.1.5	Emerald Ash Borer.....	8
610.3.1.6	Heterobasidion Root Disease.....	8
610.3.2	Funding.....	9
610.3.3	Special Projects.....	9

610.4	DEER BROWSE.....	9
610.5	INVASIVE SPECIES.....	10
610.5.1	Funding and Partnerships.....	10
610.5.2	Best Management Practices.....	11
610.5.3	Current Plant Invasive.....	11
610.5.3.1	Buckthorn.....	11
610.5.3.2	Garlic Mustard.....	12
610.5.3.3	Honeysuckle.....	12
610.5.3.4	Wild Chervil.....	13
610.5.3.5	Multiflora Rose.....	13
610.5.3.6	Reed Canary Grass.....	14
610.5.3.7	Hemp Nettle.....	14
610.5.3.8	Common Tansy.....	14

## **600 PROTECTION**

### **OBJECTIVE**

To protect and manage the resources of the County Forest from preventable losses resulting from fire, insects, diseases and other destructive elements, including those caused by people. Protective methods shall include proper silvicultural methods.

The DNR provides statewide technical guidance that will be used to inform local decisions. This guidance will be referenced to make decisions at the County level.

## **605 FIRE CONTROL**

Damage to the County Forest caused by uncontrolled fire can create an important challenge in the management of the County Forest. Loss of resource values caused by fire will be minimized through organized prevention, detection and suppression methods. Maintaining a healthy County Forest is key to fire management. The DNR is responsible for all matters relating to the prevention, detection and suppression of forest fires outside the limits of incorporated villages and cities, as stated in s. 26.11(1), Wis. Stats. The DNR works cooperatively with local fire departments in all fire control efforts. The majority of the Chippewa County Forest is part of the Intensive Forest Fire protection area. The *Fire Management Handbook*, (No. 4325.1) shall serve as the guidelines for fire control activities. The Chippewa County Department of Land Conservation & Forest Management agrees to follow WI DNR fire control guidance on the prevention and control of forest fires. If conditions are such that trails need to be closed, Chippewa County will temporarily shut down all trail activity.

### **605.1 COOPERATION WITH THE DEPARTMENT OF NATURAL RESOURCES**

Pursuant to s. 26.11(4) and s. 28.11(4)(f), Wis. Stats., and of the Chippewa County Forest Lands Ordinance, Sec. 16-4(7), the County may cooperate with the DNR in the interest of fire prevention, detection and suppression on the County Forest. This is accomplished through agreements authorizing the DNR to use County Forest land, or to utilize County personnel and equipment for fire protection activities.

#### 605.1.1 Personnel

Forest & Trails Division personnel, upon request from the DNR, shall be made available for forest fire control efforts within the County in accordance with an established Memorandum of Understanding (MOU). The DNR is responsible for training and directing the activities of County personnel in accordance with the rules identified in the *Fire Management Handbook*, (No. 4325.1).

#### 605.1.2 Equipment

Forest & Trails Division equipment, upon request and as identified in the MOU, shall be available for forest fire control suppression. During periods of high fire hazard, all Forest & Trail Division vehicles and/or crews should be equipped with one or more backpack cans, axes or shovels, appropriate personal protective equipment, mobile communication and any other equipment deemed essential by the MOU. All hand tools shall be maintained and provided by the DNR.

#### 605.1.3 Fire Detection

Fire detection is the responsibility of the DNR. Forest & Trails Division personnel may assist and report any wildfires to the DNR or 911 Dispatch.

#### 605.1.4 Forest Fire Prevention

DNR fire control personnel are authorized by the County to place fire prevention signs at recreational areas and other strategic locations within the County Forest. The County conducts and controls all operations (including harvesting) in the County Forest in a manner designed to prevent forest fires. The Forest & Trails Division and the Department personnel will coordinate during high fire danger periods to impose any necessary restrictions. These restrictions may include, but are not limited to, recreation and logging.

### 605.2 DEBRIS BURNING

Unauthorized burning of debris will not be permitted on County Forest lands, pursuant to s. 26.12(5), Wis. Stats.

### 605.3 CAMPFIRES

During periods of high fire danger, use of campfires may be restricted.

### 605.4 PRESCRIBED BURNING

All prescribed burning on County Forest lands will follow the DNR recommendations. See *Prescribed Burn Handbook* (No. 4360.5) for details. Prescribed fire may be an effective management tool in the County Forest.

### 605.5 COUNTY FOREST FIRE HAZARD AREAS

The DNR places primary emphasis on fire control efforts in pine areas. Maps of these areas are available at the local DNR field office. The County will cooperate with DNR Fire Control in providing for firebreaks or access ways. Existing access roads, firebreaks and water access points will be maintained, as deemed necessary. Secondary emphasis will be placed on hardwood areas with no firebreaks developed or maintained, however, access roads will be maintained as defined in Chapter 700 of this plan.

## 610 CONTROL OF FOREST PESTS & PATHOGENS

### 610.1 DETECTION

Damage to the County Forest caused by insects, other pests and diseases can adversely affect management of the forest resources. Losses to resource values impacted by forest pests will be minimized through integrated pest management methods, with emphasis on silvicultural prescriptions (timber sales). The detection and control of pest problems will be accomplished by County and DNR personnel, in cooperation with other agencies.

### 610.2 PEST SURVEYS

Pest surveys are conducted under the direction of the DNR's forest health specialists. The County may cooperate by providing personnel and equipment to assist in these operations.

### 610.3 SPECIFIC PESTS AND PATHOGENS OF CONCERN

Integrated pest management for the purpose of this Plan, is defined as follows:

*“The maintenance of destructive agents, including insects, at tolerable levels, by the planned use of a variety of preventive, suppressive, or regulatory tactics and strategies that are ecologically and economically efficient and socially acceptable.”*

The integrated pest management control and methodology shall be determined jointly by the County Forest Administrator and DNR Liaison Forester, in consultation with the DNR District Forest Health Specialist. Suppression of forest pests may include the following:

1. Silvicultural prescriptions, including timber sales.
2. Biological control.
3. Chemical control.

#### 610.3.1 Specific Pests of Interest

##### 610.3.1.1 Gypsy Moth

Gypsy moth (*Lymantria Dispar*) was introduced into the United States in 1869 from Europe. The caterpillars defoliate over 300 species of trees and shrubs causing some of the largest defoliation impacts in North America. Population outbreaks, approximately 10 years apart, may defoliate entire forests. The eastern 2/3<sup>rd</sup>s of Wisconsin, including Clark and Taylor Counties, is quarantined as the infestation moves westward. DATCP administers the quarantine and aerial spray suppression efforts in Wisconsin. It will be Chippewa County’s strategy to employ sound silvicultural practices to combat this pest.

##### 610.3.1.2 Oak Wilt

Oak wilt (*Bretziella fagacearum*) is a destructive disease of oak trees. It is responsible for the death of thousands of oak trees in forests, woodlots, and home landscapes each year. Oak wilt is caused by a fungus that invades and impairs the tree's water conducting system, resulting in branch wilting and tree death. Trees in both the red oak group and white oak group are affected. There is no known cure once a tree has oak wilt. Prevention of new oak wilt infection centers on the best management option and involves avoiding injury to healthy trees and removing dead or diseased trees. Counties should use the *Oak Harvesting Guidelines to Reduce the Risk of Introduction and Spread of Oak Wilt* for management guidance.

Chippewa County's strategy will be to prohibit timber harvests in red oak dominant stands between April 1 and July 15. The County may choose to remove infected trees and debark or chip them when located in a high intensive use area. Once chipped or debarked, the materials should be covered with plastic for a period of six months to kill the fungus and any insects in the material. A vibratory plow may also be used to sever root grafts on isolated pockets where feasible.

#### 610.3.1.3 Forest Tent Caterpillar

Forest tent caterpillar (*Malacosoma disstria*) can be found throughout the United States and Canada, wherever hardwoods grow. The favored hosts in Wisconsin are aspen and oak. This native insect causes region-wide outbreaks at intervals from 10 to 15 years; outbreaks usually last 2 - 5 years in the Lake States. Severe and repeated defoliation can lead to dieback and/or reduced growth of affected trees, which in some instances may be significant. Populations are often controlled by natural enemies, helping the population crash. Aerial spraying of insecticides can be an option for control as well. It will be Chippewa County's strategy to employ sound silvicultural practices to combat this cyclic pest.

#### 610.3.1.4 Two-Lined Chestnut Borer

The two-lined chestnut borer (*Agrilus bilineatus*) is a common secondary pest in trees which have been severely defoliated several years in a row. Oaks that are under stress from drought and/or defoliation by insects such as gypsy moth (*Lymantria dispar*), fall cankerworm (*Alsophila pometaria*), and forest tent caterpillar (*Malacosoma disstria*) can be infested and killed by two-lined chestnut borer. Prevention of two-lined chestnut borer through sound silvicultural practices is the best management option. Postponing management activities in stressed stands for two (2) years after severe drought and/or defoliation have ended may provide time for trees to recover and reduce their susceptibility to two-lined chestnut borer attack. Infestations should be salvaged promptly. Chippewa County will strive to maintain healthy trees through sound silvicultural practices to discourage infestation.

#### 610.3.1.5 Emerald Ash Borer

The emerald ash borer (*Agrilus planipennis*) was accidentally introduced to North America from Asia in 2002. Emerald ash borer (EAB) infestations in Wisconsin have resulted in widespread mortality to *Fraxinus* species including green, white, and black ash. It is expected that ninety-nine percent (99%) of the ash trees in Wisconsin will die.

Ash comprises a significant component in the northern hardwood timber type and can be found in nearly pure stands in some lowland areas. Adult EAB beetles feed on foliage but it is the larvae that cause mortality by feeding on the phloem and outer sapwood of the ash trees.

The *Emerald Ash Borer Silviculture Guidelines* are available to help resource managers make informed stand-level decisions to manage forests that are not yet infested by EAB, as well as implement salvage harvests and rehabilitation in stands that have already been impacted by EAB.

It is Chippewa County's strategy to utilize merchantable ash prior to passive stand conversion in compliance with the *Emerald Ash Borer Silviculture Guidelines*. Evaluation of natural regeneration following harvests will allow the County to assess the need for site preparation or artificial regeneration efforts.

#### 610.3.1.6 Heterobasidion Root Disease

Heterobasidion Root Disease (HRD, previously called annosum root rot, is caused by the fungus, *Heterobasidion irregulare*. It is a serious disease that causes pine and spruce mortality in Wisconsin, but over 200 woody species have been reported as hosts. Red and white pine trees are most commonly affected in plantation-grown stands subjected to thinning. The disease was first confirmed in Wisconsin in 1993 and has since been found in a number of counties throughout Wisconsin. Diseased trees, including overstory trees and understory seedlings and saplings, will show fading, thin crowns with tufted foliage, and eventual mortality. Currently there are no curative treatments to eliminate the HRD pathogen from a stand once it is infested, so preventing disease introduction is the best approach.



Infection most often occurs when HRD spores land and germinate on a freshly cut stump. The pathogen then grows into the root tissue and progresses underground from tree to tree through root contact. As the pathogen spreads, and trees decline and die, an ever-expanding pocket of mortality is formed. HRD fruit bodies, or conks, may be found at the base of dead trees and old stumps. Fruit bodies are most commonly observed in the fall but can be found any time of the year.

It is Chippewa County's strategy to follow the *Guidelines for stump treatment to reduce the risk of introduction and spread of Heterobasidion root disease in Wisconsin*. The HRD guidelines are designed to help property managers and landowners determine whether the preventive pesticide treatment should be used to reduce the risk of introduction and spread of HRD at the time of harvest in a pine and/or spruce stand.

#### 610.3.2 Funding

The County Forest will make all reasonable efforts to secure funding for control efforts, through County funds, or other state, federal or private funding sources.

#### 610.3.3 Special Projects

The County may cooperate with other agencies in forest pest research.

### 610.4 DEER BROWSE

Forest regeneration and reproduction is critical to sustain both timber production and wildlife habitat, and the overall health of the deer herd. As a keystone species, deer can affect forest regeneration, long-term forest production and forest sustainability. This is a concern for all interested in forest production and trying to balance deer numbers with habitat. Chippewa County Forest may monitor herbivory impacts during forest reconnaissance or build exclosures as a demonstration.

## 610.5 INVASIVE PLANT SPECIES

Invasive plants can cause significant negative impacts to the County Forest. Invasive species can displace native plants and hinder the forest regeneration efforts. Preventing them from dominating habitats is critical to the long-term health of the County Forest. There are a number of invasive plant species in varying densities in the County Forest. Some warrant immediate and continual treatment efforts while others may be allowed to remain due to extent and financial ability to control them. The County will continue to train staff in invasive species identification, as well as attempt to secure funding sources to control them, as much as is practical. Invasive plants in the County Forest should be documented, as well as potential response to new infestations.

### 610.5.1 Funding and Partnerships

Grant opportunities for invasive species control funding can be found on the “Financial Assistance” webpage of the *Wisconsin Invasive Species Council*. The number of grants for local governments and County Forest is limited, especially for terrestrial invasive plant control. Some grants, such as the Department of Natural Resource’s *Wild Turkey Stamp Program*, support invasive plant control as part of larger efforts to promote certain outcomes and might be applicable.

The Department of Natural Resources promotes the formation of Cooperative Invasive Species Management Areas (CISMAs) through its *Weed Management Area – Private Forest Grant Program*. While activities funded by this grant are restricted to non-industrial private forests, CISMAs are encouraged to partner with other groups in their area, and some can provide technical support to County Forests. The CISMA’s of Wisconsin can be found at [www.ipaw.org](http://www.ipaw.org). The County works with the Lower Chippewa Invasive Partnership, Inc. to improve monitoring for invasive in the region.

### 610.5.2 Best Management Practices

In 2009, the Department of Natural Resources and many stakeholder groups approved a series of Best Management Practices (BMPs) for minimizing the spread of forest invasive plants. The full text of the [BMPs](#) is found on the *Wisconsin Council on Forestry* website. Voluntary use of the BMPs during forestry stewardship activities reduces the spread of invasive plants that can impede forest regeneration in county forests.

BMPs used before, during and after a harvest promote forest regeneration. Reasonable efforts to clean vehicles, equipment, footwear and other clothing helps reduce the spread of seeds and plant fragments to un-infested forests. Planning the sequence and timing of stewardship activities to reduce contact with invasive plants during forestry operations is another helpful strategy. Similarly, controlling populations of invasive plants before logging reduces the risk of spreading them. Follow-up monitoring of disturbed stands can detect populations of invasive plants while they are still small and more easily managed.

### 610.5.3 Current Plant Invasives

#### 610.5.3.1 Buckthorn

Two species of invasive buckthorn impact Wisconsin's forests. Common buckthorn (*Frangula cathartica*) is more often found growing on well-drained soils while glossy buckthorn (*Frangula alnus*) favors wetter soils. Both species grow in shade or sun, quickly form dense, even-aged thickets that shade out understory plants, including tree seedlings, and hinder forest regeneration. Their dark colored fruits are eaten by birds who disperse them long distances. Both buckthorns green-up before native plants and remain green after the natives drop their leaves. Buckthorn can be controlled by taking advantage of the longer period in which they retain their leaves. Foliar applications of herbicide applied when buckthorn has leaves and the natives are leafless will minimize damage to native plants. Other control options include mowing the shrubs and then treating re-sprouts with foliar herbicide, basal bark herbicide applications, and cut stump herbicide applications.

#### 610.5.3.2 Garlic Mustard

Garlic mustard (*Alliaria petiolata*) is an herbaceous biennial, native to Europe. During the first year, a basal rosette of only leaves develops. The second year, several stems from 1 – 4 feet tall grow from the basal rosette. The leaves have a distinct garlic fragrance when crushed. From the stems grow several small white flowers. Each plant can produce 100's of tiny seeds inside long, narrow capsules. Garlic mustard can quickly colonize disturbed forests as it often follows corridors, such as game trails or man-made roads/paths. As garlic mustard spreads, it quickly displaces native plants and is known to radiate chemicals into the soil that disrupt associations between mycorrhizal fungi and native plants. Small populations can be hand pulled, while larger populations are better controlled with prescribed fire and/or herbicide. All pulled plant materials should be bagged and removed from the forest as seeds have been known to mature on dead plants left on site. Treatment should be repeated until the seed bank is depleted, which takes multiple years. Garlic mustard sites should be monitored annually, until no plants are discovered for several years.

#### 610.5.3.3 Honeysuckle

Bush Honeysuckles (*Lonicera maackii*, *L. tatarica*, *L. morrowii*, *L. X bella*) were introduced from Eurasia as ornamentals, wildlife cover and soil erosion control. Bush honeysuckles are upright deciduous shrubs, ranging from 5 - 12 feet tall with gray shaggy bark. The leaves are opposite, simple, oval and untoothed and can be smooth to velvety, depending on species. Flowers are fragrant and tubular ranging in colors of white, red and pink. They bloom May through June, and then form red to yellow berries that are found as pairs on the leaf axils. Honeysuckles replace native forest shrubs and herbaceous plants by inhibiting growth of understory plants due to early leaf-out which shades out herbaceous ground cover and depletes soil moisture. Control options include hand pulling small infestations and prescribed burning, which kills seedlings and top kills mature shrubs. Herbicide options include cut stump treatment and foliar spraying. With all control efforts, repeated monitoring is needed.

#### 610.5.3.4 Wild Chervil

Wild Chervil (*Anthriscus sylvestris*) is an herbaceous monocarpic perennial, native to Europe, with grooved hollow stems covered in soft hairs and alternate fern-like leaflets with the leaf base clasping the stem. It can grow 3 – 6 feet tall with white umbel flowers that bloom in late May-July, producing 2 long-ribbed brown seeds with narrow antenna-like tips. It grows in a variety of soil types though prefers rich, moist soils. Commonly found along roadsides and disturbed areas, it invades forests, fields and pastures. Small populations can be hand pulled provided the entire tap root is removed. Gloves, long sleeves and pants should be worn when handling this plant as it may cause skin irritation. Repeated mowing can prevent seed set. Chemical control should be applied prior to blooming.

#### 610.5.3.5 Multiflora Rose

Multiflora Rose (*Rosa multiflora*) is a thorny, perennial shrub with arching stems and leaves divided into 5-11 sharply toothed leaflets native to Asia. It forms thickets that can reach 10-15 feet tall and 9-13 feet wide. It can tolerate a wide range of soil and environmental conditions, though does best on well-drained soils. It is extremely prolific and invades roadsides, woodlands, fields, and prairies. Mechanical control can exhaust the root stock by mowing 3-6 times a year for 2-4 years. Pulling may not be effective as resprouting may occur if the entire root system is not removed. Control by herbicides with a cut stump treatment or foliar application of glyphosate or imazapyr. Basal bark treatment with triclopyr has also been effective.

#### 610.5.3.6 Reed Canary Grass

Reed Canary Grass (*Phalaris arundinacea*) is a cool-season perennial found along lakes, streams, wetlands and in moist, open areas. It can grow 2-9 feet tall with erect, hairless stems. Densely clustered single florets, green to purple when in bloom from May to mid-June, turn tan as seeds form. It dominates a significant number of wetlands in the Midwest and is widely established in the County Forest and across the State. Complete eradication is unlikely and control efforts should focus on preventing new infestations.

#### 610.5.3.7 Hemp Nettle

Hemp Nettle (*Galeopsis tetrahit*) is an herbaceous annual from Eurasia which grows 1-3 feet tall, with a square stem, swollen at leaf nodes, covered in downward pointing hairs intermixed with shorter granular hairs. Purple to pink to white flowers in dense axillary clusters with 5 sharp protruding points bloom from June-September. It prefers disturbed sites, invading roadsides, woods, pastures and fields. Minimizing the spread depends on cleaning logging equipment, ATV's, vehicles, shoes and clothing after entering infested areas. Chemical control may also be achieved using a mixture of Dicamba (230 ml/ha) with MCPA (1.1L/ha).

#### 610.5.3.8 Common Tansy

Common Tansy (*Tanacetum vulgare*) is a perennial herbaceous plant that grows 2-5 feet tall, unbranched except for the bright yellow flower head which blooms from July-October. The stem is slightly hairy, woody and purplish red near the ground. It prefers well-drained or sandy soils in open disturbed areas, roadsides, fields, prairies and pastures. Mechanical treatment by mowing prior to flowering prevents seed set. Chemical control by spraying the rosettes in the spring using Dicamba, glyphosate, or clopyralid mixed with a surfactant is also effective.